



COL17A1 gene

collagen type XVII alpha 1 chain

Normal Function

The *COL17A1* gene provides instructions for making a protein that is used to assemble type XVII collagen. Collagens are a family of proteins that strengthen and support connective tissues, such as skin, bone, tendons, and ligaments, throughout the body. In particular, type XVII collagen plays an essential role in strengthening and stabilizing the skin.

The protein produced from the *COL17A1* gene is known as a pro- α 1(XVII) chain. Three identical pro- α 1(XVII) chains twist together to form a triple-stranded, ropelike molecule known as a procollagen. Procollagen molecules are secreted by the cell and processed by enzymes to remove extra protein segments from the ends. Once these molecules are processed, they arrange themselves into long, thin bundles of mature type XVII collagen.

Type XVII collagen is a major component of hemidesmosomes, which are microscopic structures on the inner surface of the top layer of skin (the epidermis). These structures help to anchor the epidermis to underlying layers of skin. Type XVII collagen is critical for the stability of hemidesmosomes, and therefore it plays an important role in holding the layers of skin together.

Health Conditions Related to Genetic Changes

junctional epidermolysis bullosa

More than 60 mutations in the *COL17A1* gene have been identified in people with junctional epidermolysis bullosa (JEB). Most of these mutations insert or delete several DNA building blocks (base pairs) in the *COL17A1* gene or create a premature stop signal in the instructions for making the pro- α 1(XVII) chain. These changes reduce the amount of functional type XVII collagen in the skin. Without enough of this collagen, the epidermis is only weakly attached to underlying layers of skin. Friction or other minor trauma (such as rubbing or scratching) can cause the skin layers to separate, leading to the formation of blisters.


Most *COL17A1* gene mutations cause the milder form of junctional epidermolysis bullosa, known as non-Herlitz JEB. Affected individuals experience blistering, but it may be limited to the hands, feet, knees, and elbows and often improves after the newborn period. A few individuals with mutations in the *COL17A1* gene have had

the more severe form of the disorder, Herlitz JEB. Infants with Herlitz JEB develop widespread blistering that causes life-threatening complications.

Chromosomal Location

Cytogenetic Location: 10q25.1, which is the long (q) arm of chromosome 10 at position 25.1

Molecular Location: base pairs 104,031,288 to 104,085,880 on chromosome 10 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- alpha 1 type XVII collagen
- BA16H23.2
- BP180
- BPAG2
- bullous pemphigoid antigen 2 (180kD)
- COHA1_HUMAN
- collagen type XVII alpha 1
- collagen XVII, alpha-1 polypeptide
- collagen, type XVII, alpha 1
- FLJ60881
- KIAA0204
- LAD-1

Additional Information & Resources

Educational Resources

- Molecular Biology of the Cell (fourth edition, 2002): Collagens Are the Major Proteins of the Extracellular Matrix
<https://www.ncbi.nlm.nih.gov/books/NBK26810/#A3551>

GeneReviews

- Junctional Epidermolysis Bullosa
<https://www.ncbi.nlm.nih.gov/books/NBK1125>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28COL17A1%5BTIAB%5D%29+OR+%28%28type+XVII+collagen+%5Btiab%5D+AND+alpha+1+%5Btiab%5D%29+OR+%28BP180%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1440+days%22%5Bdp%5D>

OMIM

- COLLAGEN, TYPE XVII, ALPHA-1
<http://omim.org/entry/113811>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_COL17A1.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=COL17A1%5Bgene%5D>
- HGNC Gene Family: Collagens
<http://www.genenames.org/cgi-bin/genefamilies/set/490>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=2194
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/1308>
- UniProt
<http://www.uniprot.org/uniprot/Q9UMD9>

Sources for This Summary

- Bauer JW, Lanschuetzer C. Type XVII collagen gene mutations in junctional epidermolysis bullosa and prospects for gene therapy. *Clin Exp Dermatol*. 2003 Jan;28(1):53-60. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12558632>
- Franzke CW, Has C, Schulte C, Huilaja L, Tasanen K, Aumailley M, Bruckner-Tuderman L. C-terminal truncation impairs glycosylation of transmembrane collagen XVII and leads to intracellular accumulation. *J Biol Chem*. 2006 Oct 6;281(40):30260-8. Epub 2006 Aug 9.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16899459>
- Fu CL, Giudice GJ, Van den Bergh F. Protein structural analysis of BP180 mutant isoforms linked to non-Herlitz junctional epidermolysis bullosa. *J Invest Dermatol*. 2006 Jan;126(1):232-4.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16417243>
- Gatalica B, Pulkkinen L, Li K, Kuokkanen K, Ryyänen M, McGrath JA, Uitto J. Cloning of the human type XVII collagen gene (COL17A1), and detection of novel mutations in generalized atrophic benign epidermolysis bullosa. *Am J Hum Genet*. 1997 Feb;60(2):352-65.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/9012408>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1712405/>
- Nakamura H, Sawamura D, Goto M, Nakamura H, Kida M, Ariga T, Sakiyama Y, Tomizawa K, Mitsui H, Tamaki K, Shimizu H. Analysis of the COL17A1 in non-Herlitz junctional epidermolysis bullosa and amelogenesis imperfecta. *Int J Mol Med*. 2006 Aug;18(2):333-7.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16820943>
- Pasmooij AM, Pas HH, Jansen GH, Lemmink HH, Jonkman MF. Localized and generalized forms of blistering in junctional epidermolysis bullosa due to COL17A1 mutations in the Netherlands. *Br J Dermatol*. 2007 May;156(5):861-70. Epub 2007 Jan 30.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17263807>
- Pulkkinen L, Uitto J. Mutation analysis and molecular genetics of epidermolysis bullosa. *Matrix Biol*. 1999 Feb;18(1):29-42. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10367729>
- Ruzzi L, Pas H, Posteraro P, Mazzanti C, Didona B, Owaribe K, Meneguzzi G, Zambruno G, Castiglia D, D'Alessio M. A homozygous nonsense mutation in type XVII collagen gene (COL17A1) uncovers an alternatively spliced mRNA accounting for an unusually mild form of non-Herlitz junctional epidermolysis bullosa. *J Invest Dermatol*. 2001 Jan;116(1):182-7.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11168815>
- Varki R, Sadowski S, Pfindner E, Uitto J. Epidermolysis bullosa. I. Molecular genetics of the junctional and hemidesmosomal variants. *J Med Genet*. 2006 Aug;43(8):641-52. Epub 2006 Feb 10.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16473856>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2564586/>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/COL17A1>

Reviewed: September 2009
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services